



ocean periodontal

Tel: (902) 446-2326

Fax: (902) 446-2320

Email: perio@oceanperiodontal.com

www.oceanperiodontal.com

DR. CHRIS CAMERON

DR. ADITYA PATEL

DR. NADER HAMDAN

DR. ANIL SACHDEV

DR. KUDIRAT JIMOH

Halifax Professional Centre

1000-5991 Spring Garden Road

Halifax, NS B3H 1Y6

PERIODONTAL POST-SURGICAL CARE INSTRUCTIONS

PERIODONTICS
IMPLANT SURGERY

Please follow these suggestions in caring for your mouth after periodontal surgery. They are necessary for both comfort and to obtain good results.

- General**
- **NO SMOKING FOR 2 weeks**—smoking prevents good healing.
 - Limit strenuous physical activity for the next few days. You may, however, participate in normal daily activities.
 - Avoid the following for the first 48 hours: a) alcohol, b) caffeine (colas, teas, coffee), c) using straws, d) spitting, and e) blowing your nose excessively.

- Bleeding**
- For the first 24 hours, some oozing and discoloration of saliva is normal.
 - Keep your head elevated during sleep/rest.
 - For excessive bleeding and continuous flow, utilize a moist gauze or black tea bag (moistened with cold water) and apply **FIRM** pressure at bleeding site for 30 seconds. Try this at least twice.
- If bleeding does not stop, please call the After hours phone number listed below

- Swelling**
- Keep ice bag or a frozen bag of vegetables over the face, close to surgery area for 24 hours, alternating 20 minutes on and 20 minutes off.
- Swelling will be the greatest in the next 3-5 days
- If swelling and/or discomfort persists after day 5 or it seems to be getting worse, please call the After hours number below.

- Medication**
- Take the prescribed medication before the local anesthetic (freezing) wears off. Do not mix your medication with alcohol.
- Ibuprofen (Advil) 400mg OR
 - Acetaminophen (Tylenol) 650mg
- Take the above dosage every 4-6 hours for the first 2 days after surgery, and then as needed for pain
- Other medication:** _____
- Antibiotic:** _____

Diet

Eat soft, warm, or cold foods for 1-2 weeks. Examples: ice cream, milk shakes, soups, mashed potatoes, scrambled eggs, pasta, puddings, yogurts, instant breakfast, etc. and progress as tolerated to a normal diet. Maintain a good fluid intake.

- Mouth care**
- Avoid all rinsing the mouth for the first 12 hours following surgery (unless specified otherwise).
- Avoid brushing/flossing in the surgical area for 1-2 weeks while the sutures are intact. In this time, you can continue brushing/flossing all other areas as normal.
- Periodontal pack/dressing may be put on the surgical site. These dressings may fall off by themselves in pieces, or completely the next day or last till your post-op appointment. There is no set time for them to stay on.
 - Rinse with either salt water chlorhexidine solution at least twice a day for at least 30 secs. Rinsing will help keep plaque/food away from the surgical site.
- Please do not attempt to cut stitches as this may affect the outcome of the surgery.

- Sinus precaution**
- In the event of sinus grafting/involvement, please follow the instructions below:
- Avoid blowing through the nose
 - Keep mouth open during sneeze
 - Do not bend head down and keep head elevated during sleep/rest
 - Utilize over the counter decongestants such as Sudafed to keep the nasal draining continuous
 - Avoid flying for at least 3 weeks.

Additional notes

AFTER HOURS PHONE NUMBERS

- DR. CHRIS CAMERON902-880-9198
- DR. ADITYA PATEL 902-403-6032
- DR. NADER HAMDAN204-952-5717
- DR. ANIL SACHDEV902-457-2645
- DR. KUDIRAT JIMOH902-322-9696